







U.S. Army Corps of Engineers (33 CFR:1984)

"Jurisdictional wetlands"

"Those areas that are inundated or saturated by surface or ground water at a frequency and duration sufficient to support, and that under normal circumstances do support, a prevalence of vegetation typically adapted for life in saturated soil conditions. Wetlands generally include swamps, marshes, bogs, and similar areas."

Riparian Definitions

"associated with water courses. Riparian may refer to vegetation associated with large rivers or with small, even intermittent drainages such as arroyos."--Dick Peddie and Hubbard (1977)

"Riparian areas are three-dimensional ecotones of interaction that include terrestrial and aquatic ecosystems, that extend down into the groundwater, up above the canopy, outward across the floodplain, up the near-slopes that drain to the water, laterally into the terrestrial ecosystem, and along the water course at a variable width"

-- Ilhardt et al. (2000)

"a distinct ecological site, or combination of sites, in which soil moisture is sufficiently in excess of that otherwise available locally, due to run-off and/or subsurface seepage, so as to result in an existing or potential soil-vegetation complex that depicts the influence of that extra soil moisture."


-- Anderson (1987)

"environs of freshwater bodies, watercourses, and surface-emergent aquifers (springs, seeps, and oases) whose transported waters provide soil moisture in excess of that otherwise available through local precipitation to potentially support the growth of mesic vegetation."

--Warner and Hendrix (1984)

"Riparian areas are transitional between terrestrial and aquatic ecosystems and are distinguished by gradients in biophysical conditions, ecological processes, and biota. They are areas through which surface and subsurface hydrology connect waterbodies with their adjacent uplands. They include those portions of terrestrial ecosystems that significantly influence exchanges of energy and matter with aquatic ecosystems (i.e., a zone of influence). Riparian areas are adjacent to perennial, intermittent, and ephemeral streams, lakes, and estuarine-marine shorelines."

-- National Research Council (2002)



"Riparian areas generally do not satisfy regulatory and other definitions of "wetland," and thus are not encompassed by regulatory programs for wetland protection."

National Research Council, 2002











Benefits and Functions

- Intercepts sediment, nutrients, pesticides and other materials in surface runoff
- Reduces nutrients and other pollutants in shallow subsurface water flow
- Provides food and cover for wildlife including beneficial insects
- Lowers water temperatures by shading
- Slows out-of-bank flood flows


Benefits and Functions - con't ...

- Provides litter fall
 Diversifies farm and large woody debris needed by aquatic organisms
- Tree and shrub roots increase bank and shoreline stability
- enterprise timber, wood fiber, horticultural products
- Improves aesthetics, land values, reduces taxes in some locales

DEFINITIONS

Buffer Strip - "An area of vegetation adjacent to a water body that is managed to maintain the integrity of stream channels and shorelines;

of upland sources of pollution by trapping, filtering, and converting sediments, nutrients, and other chemicals."









USDA Riparian Forest Buffer

Conservation Practice Job Sheet

391

Natural Resources Conservation Service (NRCS)

April 1997

Landowner_







Figure 2.40: The width of the vegetation buffer influences filter and barrier functions.

Dissolved substances, such as nitrogen, phosphorus, and other nutrients, entering a vegetated stream corridor are restricted from entering the channel by friction, root absorption, clay, and soil organic matter.

Adapted from Ecology of Greenways: Design and Function of Linear Conservation Areas. Edited by Smith and Hellmund. © University of Minnesota Press 1993.



Purpose: To encourage the use of conservation buffer strips by agricultural producers and other landowners-- in both urban and rural settings



Goal: To install 2 million miles (up to 7 million acres) of conservation buffer strips by the year 2002



CONSERVATION BUFFERS WORK...













Army Corps of Engineers' Ecosystem Restoration Authority

ER 1105-2-100 Corps Planning Guidance 22 April 2000

"Ecosystem Restoration is one of the primary missions of the Corps of Engineers Civil Works program. The Corps objective in ecosystem restoration is to contribute to national ecosystem restoration."

"Those restoration opportunities that are associated with wetlands, riparian and other floodplain and aquatic systems are most appropriate for Corps involvement."



Water Resources Development Acts

• Authorize the Secretary of the Army to study and/or implement various projects and programs for improvements to rivers and harbors of the United States.



Corps Study Authorities To Support Ecosystem Restoration

- Continuing Authorities Program (CAP)
 - Section 1135 Ecosystem Restoration
 - Section 206 Aquatic Ecosystem Restoration
- General Investigations (GI)
 - Projects Authorized By Congress Through Passage of Resolution or Act



Objective of the Section 1135 Program

◆ The Objective of the Section 1135 Program is to restore structure, function, or dynamic processes of degraded ecosystems.

The focus in ecosystem restoration is on those ecological resources and processes that are directly associated or directly dependent upon the hydrologic regime of the ecosystem and watershed.



Criteria for Qualification as a Section 1135 Project

- The habitat restoration must be linked to a Corps of Engineers built project.
- Budgetary priority given where Corps projects contributed to the degradation of the ecosystem
- ◆ The restoration benefits must outweigh the project costs and be associated with fish and wildlife resources.



The Section 206 Program Aquatic Ecosystem Restoration

- Study Funding Level and Processes Identical to Section 1135 Program
- Is Not Restricted to Impacts Resulting from Corps Projects




General Investigation (GI) Studies

- GI Studies are Typically Watershed in Scale and are Initiated Where the Public Perceives a Problem and Seeks Congressional Assistance
- The Study Gains Congressional Approval and Funding
 - Senate Committee on Environment and Public Works
 - House Committee on Transportation and Infrastructure
- Congress Passes Resolution and/or Act
- Corps Receives Funds To Initiate a Study








Denton County, Texas





GREENBELT CORRIDOR
RAY ROBERTS/LEWISVILLE LAKE
FM 455 ACCESS AREA
TEXAS PARKS AND WILDLIFE DEPARTMENT










Rio Salado - (Salt River) Riparian Restoration Project















Riparian Definitions

Regulatory Program of the US Army Corps of Engineers Part 330 - Nationwide Permit Program

Vegetated buffer - A vegetated upland or wetland area next to rivers, streams, lakes, or other open waters which separates the open water from developed areas, including agricultural land. Vegetated buffers provide a variety of aquatic habitat functions and values (e.g., aquatic habitat for fish and other aquatic organisms, moderation of water temperature changes, and detritus for aquatic food webs) and help improve or maintain local water quality. A vegetated buffer can be established by maintaining an existing vegetated area or planting native trees, shrubs, and herbaceous plants on land next to open waters. Mowed lawns are not considered vegetated buffers because they provide little or no aquatic habitat functions and values. The establishment and maintenance of vegetated buffers is a method of compensatory mitigation that can be used in conjunction with the restoration, creation, enhancement, or preservation of aquatic habitats to ensure that activities authorized by NWPs result in minimal adverse effects to the aquatic environment."


Regulatory Program of the US Army Corps of Engineers

Since passage and subsequent amendments to the Clean Water Act, the Corps has had legal authority during Section 404 permit decisions to require vegetated buffer strips as part of the mitigation for filling wetlands. Both upland and riparian vegetated buffer strips can be mandated, in certain situations, by District Engineers as part of a Section 404 permit under the NWP system.

Vegetated buffer strips, such as forested riparian areas, are well-suited for protecting water quality and providing wildlife habitat, and "are a critical element of the overall aquatic ecosystem in virtually all watersheds" (Federal Reg. 67(10), p. 2064).

The Corps has also "believes we need to protect open waters better than we have in the past, and vegetated buffers are a critical element of that protection."

The Corps "believes that establishing or maintaining existing vegetated buffers to open waters is critical to overall protection of the nation's






















Riparian zones provide migration corridors for birds and many large mammals that use different summer and winter ranges.





Products



ERDCEL TN-EMPIRP-8108 January 2000

Width of Riparian Zones for Birds

BACKGROUND..Over the past several decades, more than 450 Corps of Engineers Civil. Works reservoir projects have been constructed in 43 states encompassing nearly 12 million acros (at normal pool elevations, about one half is water and the remaining half is associated

land). The majority of inland Civil Works projects are constructed along streams and rivers. There is increasing interest in managing the riperion buffer strips (i.e., vegetation adjacont to streams, rivers, and lakes) along these watercourses. Retaining riporion vegetation of proper width not only minimizes the impacts of crossion and nonpoint-source pollution; these areas also provide habitat and movement. corridors for wildlife as well as benefits to fish populations (Fischer et al. 1999) (Figure 1). Unfortunately, when decisions are made to nostore or manage buffer strips adjacent to streams and rivers, the basis for determining strip recommen contains to a variety width has been almost completely dominated by



water quality considerations. Few studies have addressed the compatibility of a buffer strip widths with other important ecological functions, especially their a native found and floral species.

Many riparian zones in North America are degraded to the point that they do no resources needed to make them suitable as habitat or as movement corridors. I also negatively affects many of the other important functions and values these provide. Providing bird habitat othen is a management objective on Corps land of riparian zones is a viable option. However, there is little guidance available management of riparian zones for birds.

IMPORTANCE TO AVIAN COMMUNITIES

About half of the approximately 720 birds that brood in North America are con nestropical migrants (Figure 2), those that broad in North America but migrate grounds in the Caribbean, Mexico, and Costral and South America. The remai resident birds (non-raignatory) or short-distance migrants (typically bread in the States and Canada and winter in the southern United States). Information from across the nation indicates that many bird species are declining, especially the micrant species.



ridors and vegetated buffer zone: delines for Corps of Engineers pr

ctively manage natural resources tendeds, it is often necessary to topotes of some reflect bottops toms from land-use activities that

ority of inland Curps of Engineers projects are constructed along.



Riparian Terminology: Confusion and Clarification

by Richard A. Fincher," Chester C. Marks." John T. Ratti," and John Guideo.



INTRODUCTION

Rowan zones occur throughout the United Status as large arrays of vacadation adjacent to streams, rivers, lakes, reservoirs, and other inland equatic systems that affect or are effected by the presence of water. This egetation contributes to unique ecosystems that perturn a large variety of ecological functions. Unfortunately, considerable variation associated with riparian familiosings, shifter to problems associated with welland terminology (Mitself- and Gossalins 1903). This our lead to confusion when people alternal is percentage about reporter pores, perfordantly If they come from different disoplinary backgrounds. The goals of this poper are to prumote asserses of this problem by describing variation associated with semantics in diselect territorings, to region why the contributes to confusion, to show the importance of alternating to standardize this terminategy, and to ouggest ways that natural resource professionals can believ describe What comprises a rowher wiresystem

WHY IS THERE CONFUSION? No Universally Accepted Ripartee Definition.

satisfy the needs of all assertiots or apercons For example, Cowardin et al. (1979)-defined welfunds and despessor habitate for the National Waltends Classification System and traentary (NWI), whereas the U. S. Army Corps of Engineers uses a different definition under Section 454 of the Clear Water Act to requise the deposition of divolged and fill materials into wellands. Similarly there is no priversally

recognized or widely accepted definition: pringularly siecel lises of Sporian Johns Anderson 1967). Rigieran definitions In earne texts are over-empiried, and books on witdiffy habitute and plant communities do not adequately disting ripariet communities from upland. communities (Chinart and Anderson)

Riparian definitions range from simple descriptions, such as "associated with water courses" (Dick-People and Hubb 1977-06), to techniqui and deballed descri-for opecific areas (e.g., Mingried et al. (19 Recently, threch et al. (2000; p. 25) proportions functional definition for openies por They suggested that riparitor across are. "I almost soutcomes of interaction that in ter within and aquatic ecosystems. But eshown into the proundwater, up above the correspy, codward across the fluodplain, up reservinges that draw to the water, lateral the terrestrial ecosystem, and along the a course at a variable width." However, the no indication that this or other recent defi-sell become universally occupied as the

Regional Differences.

Stream and river ecosystems differ region and boolly in many characteristics, inclu-width, depth, frequency of flooding. ydropeuroophic factors, and regelature hese differences are must apparent between Eastern and Western regions of the United States. Rigarian power in the and West ober copy of the order streets having extreme and sarable Resal conditions (Mitark and

*Little Reason's and Destinational Global Environmental Laboratory, 8600 rapis Funy No., 1000 rapis 500 0000.
*Company of Sales College of Fundamental Readon Statement Reasons, 200

ERIOC THEMRAP-SR-25.

Design Recommendations for Riparian Corridors and Vegetated Buffer Strips

by Richard A. Fischer⁸ and J. Craig Fischenich⁹

April 2000

Complexity

Value as a Planning Tool

INTRODUCTION

Binarian zones occur as transitional areas between aquatic and upland terrestrial habitats. Although not always well-defined (Fischer et al. 2000). They generally can be described as long. linear strips of vegetation adjacent to streams, rivers, lakes, reservoirs, and other inland aquatic systems that affect or are affected by the presence of water. Riparian zones typically comprise a small percentage of the landscape. often less than 1 percent, yet they frequently harbor a disproportionately high number of wildlife species and perform a disparate number of ecological functions when compared to most upland habitats. Riparian zones have been widely recognized as functionally unique and dynamic ecosystems only within the past 25 years. Even more recently, these areas have become a major focus in the restoration and management of landscapes (Knopf et al. 1988, Nalman, Décamps, and Pollock 1993).

Unfortunately, many riparian zones in North America do not function properly (e.g., they are degraded to the point that they do not protect water quality or provide the resources needed to make them suitable as wildlife habitat or as



Figure 1. Characteristics of vegetated riparian buffer strips influence water quality, wildlife, and recreational apportunities (photo courtesy of the U.S. Army Corps of Engineers).

movement corridors). This degradation also negatively affects many of the other important. functions and values these landscape features

WHAT IS THE DIFFERENCE BETWEEN BUFFER STRIPS AND CORRIDORS?

There is considerable confusion in the literature regarding both wetlands and riparian zones (Fischer et al. 2000). At the heart of this confusion is the proper distinction between vegetated buffer strips and corridors. Riparian zones are most commonly referred to as vegetated buffer strips (e.g., riparian buffer

ERDC TN-EMRRP-8R-24


⁶ LB Army Engineer Research and Development Center, Environmental Laboratory, 3000 mats Party Rd., Voluntury, MR 30160.

Riparian Zone Management









How Do I Design a Riparian Buffer Strip or Corridor?

FOREST BUFFER - 60'-- 20' -15'-ZONE II ZONE 1 ZONE 2 **BUNOFF CONTROL** PASTURE MANAGED FOREST UNDISTURBED MOTTOBIA Controlled Periodic harvesting is Watering Tree removal detritus for processing is generally not permitted grazing or facilities and necessary in Zone 2 to nd provide cover and haying can be permitted in livestock are remove nutrients sequesish and other stream kept out of the tered in tree stems and in this zone. Riparian Zone Zone 3 under branches and to maintain insofar as nutrient uptake through vigorous tree growth. certain conditions. practicable.

How Wide Should my Buffer Strip or Corridor Be?



Riparian Zone Widths Recommended for Water Quality Concerns

Objective	Width (m)
Reducing P Concentrations	15m
Remove 80% of excess N and P	5 m
Improve quality of stormwater runoff	25 m
Remove sediment	61 m
Remove Nitrates and Phosphorus	18 m
Reducing suspended solids	9 m

Note: These are site-specific studies and aren't intended to be general recommendations

Riparian Zone Widths Recommended for Bird Communities


Objective	State	Width (m)
Neotropical migrants	VA	50m
Neotropical migrants	KY	100m
Include 90% of bird species	$\mathbf{V}\mathbf{T}$	150m
Maintain complete avian community	\mathbf{SC}	500m
Maintain functional bird community	GA	100m
Breeding habitat for yellow-billed cuckoo	CA	100m
Reduce edge-related predation	ME	150m
Breeding habitat for area-sensitive species	NH	100m

Recommended Widths of Riparian Buffer Zones for Amphibians, Mammals, and Plants

Objective


Width (m)

Support diverse herpetofaunal community	100m
Year-round amphibian habitat	165m
General herpetofaunal habitat	30m
Maintain gray squirrel populations	50m
Include 90% of vascular plant species	30m
Maintain unaltered microclimatic gradient	45m
Maintain benthic invertebrate habitat	30m













NEWS RELEASE

For Immediate Release - July 9, 2002

CONTACTS:

Jordan Peavey, The Nature Conservancy, (703) 841-5980, <u>ipeavey@tnc.org</u> <u>http://www.nature.org</u> David Hewitt, US Army Corps of Engineers, (202) 761-0289, <u>david.w.hewitt@hq02.usace.army.mil</u>

http://www.usace.army.mil

The Nature Conservancy and U.S. Army Corps of Engineers Announce Partnership to Improve Management of U.S. Rivers



The Conservancy and the Corps have agreed on changes in the management of the Green River Dam in Kentucky. The

WASHINGTON, D.C. –The Nature Conservancy and the Army Corps of Engineers today announced a collaborative effort to improve the management of dams on various rivers across the country.

Under the new partnership, entitled the Sustainable Rivers Project, the two organizations will work together to improve dam operations, helping to restore and protect the health of rivers and surrounding natural areas while continuing to meet human needs for services such as flood control and power generation. The partnership is one that both organizations expect will improve the quality of America's waterways.

Endangered Species Act

- Riparian areas provide habitat for a large number of threatened and endangered species.
- Species such as the southwestern willow flycatcher and least Bell's vireo have been major issues on some Corps project lands.








ederally Endangered lue to large-scale loss of riparian habitat

Selected Products







NEWS RELEASE

For Immediate Release - July 9, 2002

CONTACTS:

Jordan Peavey, The Nature Conservancy, (703) 841-5980, <u>ipeavey@tnc.org</u> <u>http://www.nature.org</u> David Hewitt, US Army Corps of Engineers, (202) 761-0289, <u>david.w.hewitt@hq02.usace.army.mil</u>

http://www.usace.army.mil

The Nature Conservancy and U.S. Army Corps of Engineers Announce Partnership to Improve Management of U.S. Rivers



The Conservancy and the Corps have agreed on changes in the management of the Green River Dam in Kentucky. The

WASHINGTON, D.C. –The Nature Conservancy and the Army Corps of Engineers today announced a collaborative effort to improve the management of dams on various rivers across the country.

Under the new partnership, entitled the Sustainable Rivers Project, the two organizations will work together to improve dam operations, helping to restore and protect the health of rivers and surrounding natural areas while continuing to meet human needs for services such as flood control and power generation. The partnership is one that both organizations expect will improve the quality of America's waterways.

